

Package ‘tptest’

October 1, 2026

Type Package

Title Universal Turning Point and Inflection Point Tests

Version 1.1.0

Date 2026-09-30

Description Performs turning point and inflection point tests for U-shaped and inverse U-shaped relationships in regression models. Implements the Sasabuchi (1980) test as extended by Lind and Mehlum (2010) with support for quadratic, cubic, log-quadratic, and inverse functional forms. Features include delta-method standard errors, Fieller confidence intervals, Simonsohn (2018) two-lines test, and parametric bootstrap. Designed for post-estimation analysis of linear models, panel models, and quantile regression.
References: Lind and Mehlum (2010) <[doi:10.1111/j.1468-0084.2009.00569.x](https://doi.org/10.1111/j.1468-0084.2009.00569.x)>; Sasabuchi (1980); Fieller (1954) <[doi:10.1111/j.2517-6161.1954.tb00159.x](https://doi.org/10.1111/j.2517-6161.1954.tb00159.x)>.

License GPL-3

URL <https://github.com/muhammedalkhalaf/tptest>

BugReports <https://github.com/muhammedalkhalaf/tptest/issues>

Encoding UTF-8

LazyData true

RoxygenNote 7.3.3

Depends R (>= 4.0.0)

Imports stats, graphics

Suggests testthat (>= 3.0.0), knitr, rmarkdown, lmtest, sandwich, plm, quantreg

NeedsCompilation no

Author Muhammad Alkhalaf [aut, cre, cph] (ORCID:
<<https://orcid.org/0009-0002-2677-9246>>)

Maintainer Muhammad Alkhalaf <muhammedalkhalaf@gmail.com>

Repository CRAN

Date/Publication 2026-10-01 08:30:15 UTC

Contents

ekc	2
fieller_ci	3
tptest	5
tptest-methods	9
twolines_test	10
Index	11

ekc	<i>Environmental Kuznets Curve Data</i>
-----	---

Description

Simulated data representing the Environmental Kuznets Curve (EKC) hypothesis, which posits an inverse U-shaped relationship between environmental degradation and economic development.

Usage

ekc

Format

A data frame with 500 observations and 6 variables:

- country** Country identifier (1-50)
- year** Year (2000-2009)
- gdp** GDP per capita (thousands of dollars)
- gdp_sq** Squared GDP per capita
- emissions** CO2 emissions (tons per capita)
- emissions_log** Log of CO2 emissions

Details

This is simulated panel data designed to demonstrate the tptest package. The true data generating process follows an inverse U-shape with:

- Turning point at GDP ~ 25,000 USD per capita
- Country-specific fixed effects
- Year-specific trends

Source

Simulated data for demonstration purposes.

References

Grossman, G. M. and Krueger, A. B. (1995). Economic growth and the environment. *Quarterly Journal of Economics*, 110(2), 353-377.

Examples

```
data(ekc)
head(ekc)

# Fit quadratic model
fit <- lm(emissions ~ gdp + gdp_sq, data = ekc)
summary(fit)

# Test for inverse U-shape
result <- tptest(fit, vars = c("gdp", "gdp_sq"), data = ekc)
print(result)
```

fieller_ci

Fieller Confidence Set for the Turning Point

Description

Computes the Fieller (1954) confidence set for the turning point, which is exact for linear models and remains informative when the denominator coefficient is imprecisely estimated (Lind and Mehlum 2010, equation 8).

Usage

```
fieller_ci(b1, b2, s11, s12, s22, level = 0.95, form = "quadratic", df = NULL)
```

Arguments

b1	First coefficient (β_1)
b2	Second coefficient (β_2)
s11	Variance of b1
s12	Covariance of b1 and b2
s22	Variance of b2
level	Confidence level (two-sided). Lind and Mehlum (2010) note that the test of a U shape at level α corresponds to checking whether the $1 - 2\alpha$ interval lies inside the data range.
form	Functional form: "quadratic" ($x^* = -b_1/(2b_2)$), "inverse" ($x^* = \sqrt{b_2/b_1}$), or "logquadratic" ($x^* = \exp(-b_1/(2b_2))$).
df	Degrees of freedom for the critical value: NULL (default) uses the normal distribution, a finite number uses $t(df)$.

Details

The set for the ratio $\rho = n/d$ of two coefficients is $\{\rho : (n - \rho d)^2 \leq T^2(s_{nn} - 2\rho s_{nd} + \rho^2 s_{dd})\}$, whose boundary points are $(nd - T^2 s_{nd} \pm T\sqrt{D})/(d^2 - T^2 s_{dd})$ with $D = (s_{nd}^2 - s_{nn}s_{dd})T^2 + d^2 s_{nn} + n^2 s_{dd} - 2nds_{nd}$. For the quadratic form $\rho = b_1/b_2$ and $x^* = -\rho/2$; for the inverse form $\rho = b_2/b_1$ and $x^* = \sqrt{\rho}$, restricted to $\rho > 0$.

Value

A list with elements lo, hi and type:

"bounded" the set is the interval [lo, hi]. For the inverse and log-quadratic forms lo = 0 means the set is (0, hi].

"two_rays" the set is the union of two rays, (-Inf, lo] and [hi, Inf) ((0, lo] and [hi, Inf) for the inverse and log-quadratic forms). This happens when the denominator coefficient is not significant at level but the discriminant is positive.

"ray" the set is a half-line; one of lo, hi is infinite (or lo = 0 for the positive forms).

"unbounded" the set is the whole real line (whole positive line for the inverse and log-quadratic forms).

"empty" inverse form only: no positive turning point is compatible with the data at level.

"not_applicable" form is not supported.

References

Fieller, E. C. (1954). Some problems in interval estimation. *Journal of the Royal Statistical Society: Series B*, 16(2), 175-185. doi:10.1111/j.25176161.1954.tb00159.x

Lind, J. T. and Mehlum, H. (2010). With or without U? The appropriate test for a U-shaped relationship. *Oxford Bulletin of Economics and Statistics*, 72(1), 109-118. doi:10.1111/j.1468-0084.2009.00569.x

Examples

```
# Quadratic: b1 = -6, b2 = 0.55 with a precise b2 gives a bounded interval
fieller_ci(-6, 0.55, s11 = 0.04, s12 = -0.003, s22 = 0.0004, level = 0.95)
```

```
# Same with t(120) critical value
fieller_ci(-6, 0.55, s11 = 0.04, s12 = -0.003, s22 = 0.0004, df = 120)
```

```
# b2 not significant at 5 percent: the set is the union of two rays
fieller_ci(-2, 0.15, s11 = 0.5, s12 = -0.05, s22 = 0.01)
```

Description

Tests for U-shaped or inverse U-shaped relationships using the Sasabuchi (1980) test as extended by Lind and Mehlum (2010). Supports quadratic, inverse and log-quadratic functional forms, and a cubic form with a segment-wise extension of the test.

Usage

```
tptest(
  model = NULL,
  vars,
  coefs = NULL,
  vcov_mat = NULL,
  min = NULL,
  max = NULL,
  form = c("auto", "quadratic", "cubic", "inverse", "logquadratic"),
  level = 0.95,
  delta = TRUE,
  fieller = FALSE,
  twolines = FALSE,
  bootstrap = FALSE,
  breps = 1000,
  data = NULL,
  depvar = NULL,
  bounds_scale = c("regressor", "levels"),
  df = NULL
)
```

Arguments

model	A fitted model object (e.g., from <code>lm</code> , <code>glm</code> , <code>plm::plm</code>). Alternatively, coefficients can be provided directly via <code>coefs</code> .
vars	Character vector of length 2 or 3 giving the names of the coefficients of the regressors that carry the curvature, in this order: <code>c("x", "x_sq")</code> for the quadratic form, <code>c("x", "x_inv")</code> (the regressors x and $1/x$) for the inverse form, <code>c("lnx", "lnx_sq")</code> (the regressors $\ln x$ and $(\ln x)^2$) for the log-quadratic form, and <code>c("x", "x_sq", "x_cu")</code> for the cubic form. The first name is also used to look up the regressor column when the data range is determined automatically.
coefs	Named numeric vector of coefficients. If provided, <code>model</code> is not required. Must include names matching <code>vars</code> .
vcov_mat	Variance-covariance matrix for the coefficients. Required when <code>coefs</code> is provided.

min	Lower bound of the interval $[x_l, x_h]$ on which the test is carried out. If NULL, the minimum of the regressor named in <code>vars[1]</code> over the estimation sample is used. See <code>bounds_scale</code> for the scale of this argument.
max	Upper bound of the interval; see <code>min</code> .
form	Functional form: "auto" (default), "quadratic", "cubic", "inverse", or "logquadratic". With "auto", two names in <code>vars</code> select the quadratic form and three names select the cubic form.
level	Confidence level for intervals (default 0.95). The Sasabuchi test itself does not depend on level.
delta	Logical; compute delta-method SE and CI (default TRUE).
fieller	Logical; compute Fieller confidence set (default FALSE). Available for the quadratic, inverse and log-quadratic forms.
twolines	Logical; perform Simonsohn (2018) two-lines test (default FALSE).
bootstrap	Logical; compute parametric bootstrap CI (default FALSE).
breps	Number of bootstrap replications (default 1000).
data	Optional data frame: the data used to fit model. It is required for the two-lines test and is used as a fallback to determine the data range when the regressor cannot be recovered from the model frame (rows dropped by the model's <code>na.action</code> are then excluded).
devar	Name of dependent variable for two-lines test.
bounds_scale	Scale on which user-supplied <code>min</code> and <code>max</code> are given. "regressor" (default): the scale of the regressor named in <code>vars[1]</code> , that is x for the quadratic, inverse and cubic forms and $\ln x$ for the log-quadratic form. "levels": only meaningful for the log-quadratic form, <code>min</code> and <code>max</code> are given in levels of x and are log-transformed internally. The automatic range is always taken on the regressor scale.
df	Degrees of freedom for the t distribution used in the one-sided tests and in the critical values of the delta-method and Fieller intervals. NULL (default) selects automatically: the residual degrees of freedom of the model for <code>lm</code> -type models (including <code>plm</code>), and the normal distribution for <code>glm</code> objects, for models without <code>df.residual</code> , and for the <code>coefs</code> path. <code>Inf</code> forces the normal distribution; a finite number forces $t(df)$.

Details

Sasabuchi (1980) / Lind and Mehlum (2010) test. With $y = \beta x + \gamma f(x)$ and f' monotone on $[x_l, x_h]$, a U shape is implied by $\beta + \gamma f'(x_l) < 0 < \beta + \gamma f'(x_h)$. The null hypothesis (monotone or inverse U) is rejected at level α when both one-sided t-tests reject at level α ; the overall statistic is $\min(-t_l, t_h)$ for a U shape and $\min(t_l, -t_h)$ for an inverse U shape, and its p-value is the upper tail probability of that minimum. The alternative (U or inverse U) is chosen from the sign of the change of the slope across the interval. When the fitted extremum lies outside the interval the statistic is negative and the null hypothesis cannot be rejected; the statistic and its p-value are still reported. The reported `p_min` and `p_max` are the one-sided p-values of the two component tests under the tested alternative.

Distribution. The t distribution with the model's residual degrees of freedom is used for lm-type models. For generalized linear models the test is only asymptotically valid (Lind and Mehlum 2010, Section 2), so the normal distribution is used; the same applies to the coefs path. The same distribution is used for the delta-method and Fieller critical values; see argument df.

Functional forms:

- **Quadratic:** $y = \beta_1 x + \beta_2 x^2$; turning point at $x^* = -\beta_1/(2\beta_2)$
- **Inverse:** $y = \beta_1 x + \beta_2/x$; turning point at $x^* = \sqrt{\beta_2/\beta_1}$ (requires $\beta_2/\beta_1 > 0$; $\beta_1 > 0$ gives a U shape, $\beta_1 < 0$ an inverse U shape)
- **Log-quadratic:** $y = \beta_1 \ln x + \beta_2 (\ln x)^2$. The regressors named in vars are $\ln x$ and $(\ln x)^2$ and all computations are those of the quadratic form in $\ln x$. The turning point is reported in levels, $x^* = \exp(-\beta_1/(2\beta_2))$, with intervals transformed by exp. The bounds are on the $\ln x$ scale unless bounds_scale = "levels".
- **Cubic:** $y = \beta_1 x + \beta_2 x^2 + \beta_3 x^3$. Here f' is not monotone on an interval containing the inflection point, so the two-endpoint test of Lind and Mehlum (2010) does not apply to the whole interval (their footnote 3). As a package extension, the interval is split at the inflection point $-\beta_2/(3\beta_3)$ when it lies inside, and the two-endpoint test is applied on each sub-interval, on which the slope is monotone. The results are returned in sasabuchi\$segments. The split point is the estimated inflection point, treated as fixed, so the sub-interval tests do not have the exact size of the test in the paper; this extension is not part of Lind and Mehlum (2010). When the inflection point lies outside the interval the slope is monotone on the whole interval, the single segment is the test of equation (9) of Lind and Mehlum (2010) with $H = 3$, and t_overall and p_overall are reported; otherwise they are NA.

Value

An object of class "tptest" containing:

- tp** Turning point estimate. For the log-quadratic form it is reported in levels of x , $\exp(-b_1/(2b_2))$.
- tp_se** Delta-method standard error of tp
- tp_ci** Delta-method confidence interval for the turning point. For the log-quadratic form the interval is computed on the $\ln x$ scale and exponentiated.
- tp_log, tp_log_se** Log-quadratic form only: turning point and its delta-method SE on the $\ln x$ scale.
- shape** Description of the fitted curve on the interval: "U shape", "Inverse U shape", or a monotone label when the extremum lies outside the interval (cubic form: see Details)
- alternative** The alternative hypothesis tested by the Sasabuchi statistic ("U shape" or "Inverse U shape")
- model_form** Functional form used
- sasabuchi** List with the Sasabuchi test results: slopes and t-statistics at the bounds, one-sided p-values, the overall statistic and its p-value, and a logical outside (extremum outside the interval). For the cubic form it also contains segments.
- fieller** Fieller confidence set (if requested); see [fieller_ci](#)
- twolines** Two-lines test results (if requested)
- bootstrap** Bootstrap results (if requested)

coefficients Named vector of relevant coefficients

vcov Variance-covariance matrix

bounds Interval bounds on the regressor scale

bounds_levels Log-quadratic form only: the bounds in levels of x

df Degrees of freedom used (NULL when the normal distribution is used)

References

Lind, J. T. and Mehlum, H. (2010). With or without U? The appropriate test for a U-shaped relationship. *Oxford Bulletin of Economics and Statistics*, 72(1), 109-118. doi:[10.1111/j.1468-0084.2009.00569.x](https://doi.org/10.1111/j.1468-0084.2009.00569.x)

Sasabuchi, S. (1980). A test of a multivariate normal mean with composite hypotheses determined by linear inequalities. *Biometrika*, 67(2), 429-439.

Fieller, E. C. (1954). Some problems in interval estimation. *Journal of the Royal Statistical Society: Series B*, 16(2), 175-185. doi:[10.1111/j.25176161.1954.tb00159.x](https://doi.org/10.1111/j.25176161.1954.tb00159.x)

Simonsohn, U. (2018). Two lines: A valid alternative to the invalid testing of U-shaped relationships with quadratic regressions. *Advances in Methods and Practices in Psychological Science*, 1(4), 538-555.

Examples

```
# Simulate data with U-shaped relationship
set.seed(42)
n <- 200
x <- runif(n, 1, 10)
y <- 50 - 8*x + 0.5*x^2 + rnorm(n, sd = 5)
dat <- data.frame(y = y, x = x, x_sq = x^2)

# Fit quadratic model
fit <- lm(y ~ x + x_sq, data = dat)

# Test for U-shape (data range taken from the estimation sample)
result <- tptest(fit, vars = c("x", "x_sq"))
print(result)

# With Fieller interval and two-lines test
result2 <- tptest(fit, vars = c("x", "x_sq"),
                 fieller = TRUE, twolines = TRUE,
                 data = dat, depvar = "y")
summary(result2)
```

tpctest-methods	<i>Methods for tptest Objects</i>
-----------------	-----------------------------------

Description

Print, summary, plot, and accessor methods for tptest objects.

Usage

```
## S3 method for class 'tptest'
print(x, ...)

## S3 method for class 'tptest'
summary(object, ...)

## S3 method for class 'tptest'
plot(
  x,
  main = NULL,
  xlab = NULL,
  ylab = "Marginal Effect",
  col.line = "steelblue",
  col.tp = "red",
  col.ci = grDevices::rgb(0.2, 0.4, 0.8, 0.2),
  lwd = 2,
  n = 200,
  ...
)

## S3 method for class 'tptest'
coef(object, ...)

## S3 method for class 'tptest'
confint(object, parm = NULL, level = NULL, ...)
```

Arguments

x	A tptest object
...	Additional arguments (currently ignored)
object	A tptest object
main	Plot title
xlab	X-axis label
ylab	Y-axis label
col.line	Color for the fitted curve
col.tp	Color for the turning point marker

<code>col.ci</code>	Color for confidence band
<code>lwd</code>	Line width
<code>n</code>	Number of points for plotting curve
<code>parm</code>	Parameter specification (currently ignored)
<code>level</code>	Confidence level (default uses level from <code>tptest</code> object)

<code>twolines_test</code>	<i>Simonsohn (2018) Two-Lines Test</i>
----------------------------	--

Description

Performs the two-lines test proposed by Simonsohn (2018) as an alternative to quadratic regression for testing U-shaped relationships.

Usage

```
twolines_test(data, x_var, y_var, split_point, form = "quadratic")
```

Arguments

<code>data</code>	Data frame containing the variables
<code>x_var</code>	Name of the x variable (character). For the log-quadratic form this is the $\ln x$ column, the regressor used in the model.
<code>y_var</code>	Name of the y variable (character)
<code>split_point</code>	Point at which to split the data (usually the turning point). For the log-quadratic form it is given in levels of x , as returned by <code>tptest</code> , and the split is made at $\log(\text{split_point})$ on the $\ln x$ column.
<code>form</code>	Functional form (for log-quadratic, split is in log-space)

Value

A list with test results including slopes, t-values, p-values, and whether the test confirms the U-shape.

References

Simonsohn, U. (2018). Two lines: A valid alternative to the invalid testing of U-shaped relationships with quadratic regressions. *Advances in Methods and Practices in Psychological Science*, 1(4), 538-555.

Index

* **datasets**

ekc, [2](#)

coef.tptest (tptest-methods), [9](#)

confint.tptest (tptest-methods), [9](#)

ekc, [2](#)

fieller_ci, [3](#), [7](#)

plot.tptest (tptest-methods), [9](#)

print.tptest (tptest-methods), [9](#)

summary.tptest (tptest-methods), [9](#)

tptest, [5](#), [10](#)

tptest-methods, [9](#)

twolines_test, [10](#)